

Treatment of the Fluoroquinolone Associated Disability - the pathobiochemical implications

Krzysztof Michalak^{1,2}, Aleksandra Sobolewska- Włodarczyk³, Marcin Włodarczyk³, Justyna Sobolewska⁴, Piotr Woźniak⁴, Bogusław Sobolewski⁴

¹ Adam Mickiewicz University Poznań, Physics Faculty, Lab. of Vision Science and Optometry, Umultowska street 85, 61-614 Poznań, Poland

kmichalak@amu.edu.pl, orcid: 0000-0002-0267-5605

² Nanobiomedical Center of Poznań, Umultowska street 85, 61-614 Poznań, Poland

³ Department of Biochemistry, Medical University of Lodz, Mazowiecka street 6/8

92-215 Łódź, Poland olasobolewska1@poczta.onet.pl, dr.mwłodarczyk@gmail.com

⁴ Polish Mother's Memorial Hospital-Research Institute, Outpatient Clinic, Rzgowska street 281/289, Łódź, Poland boguslaw.sobol@op.pl, orcid: 0000-0002-6884-2330
prw.113@gmail.com, orcid: 0000-0002-7273-5618

Correspondence author:

Aleksandra Sobolewska-Włodarczyk

Department of Biochemistry, Medical University of Lodz

olasobolewska1@poczta.onet.pl

The authors declare that there is no conflict of interest regarding the publication of this paper.

Abstract

Long term Fluoroquinolone Associated Disability (FQAD) after fluoroquinolone (FQ) antibiotic therapy appears in recent years a significant medical and social problem, because patients suffer for many years after prescribed antimicrobial FQ-treatment from tiredness, concentration problems, neuropathies, tendinopathies and other symptoms. The knowledge about the molecular activity of FQs in the cells remains unclear in many details. The effective treatment of this chronic state remains difficult and not effective. The current paper reviews the pathobiochemical properties of FQs, hints the directions for further research and reviews the research concerning the proposed treatment of patients.

Based on the analysis of literature, the main directions of possible effective treatment of FQAD are proposed: (A) reduction of the oxidative stress, (B) restoring reduced mitochondria potential $\Delta\Psi_m$, (C) supplementation of uni- and bivalent cations that are chelated by FQs and probably ineffectively transported to the cell. (Caution must be paid to Fe and Cu because they may generate Fenton reaction), (D) stimulating the mitochondrial proliferation, (E) removing FQs permanently accumulated in the cells (if this phenomenon takes place), (F) regulating the disturbed gene expression and enzyme activity.

keywords:

fluoroquinolones, adverse reaction, Fluoroquinolone Associated Disability, mitochondrial potential, apoptosis, vitamin E, n-acetylcysteine, Fenton Reaction, gene expression, oxidative stress, electron transport chain, Krebs Cycle, Reactive Oxygen Species, Permeability Transition Pore, hydrogen peroxide, superoxide, hydroxyl radical, fluoroquinolone toxicity, Hypoxia Inducible Factor

1. Long Term Adverse Reactions caused by Fluoroquinolones

Fluoroquinolones (FQ) belong to the group of broad-spectrum antibiotics, effective for both gram-negative and gram-positive bacteria. The most frequently prescribed drugs are ciprofloxacin (CIP), norfloxacin (NOR) and levofloxacin (LEV). FQs employ their antibacterial effect by preventing bacterial DNA from unwinding and duplicating which takes place by inhibition of bacterial topoisomerase and gyrase. For three last decades FQs played an important role in treatment of serious bacterial infections, especially hospital-acquired infections. However, due to the possibility of serious side-effects, these drugs are not currently first-line medicines and their use becomes more restrictive and limited. FQs should be reserved for those who do not have alternative treatment options.

In 2016 the US Food and Drug Administration (FDA) updated warnings, using next “black box” for oral and injectable FQs. The authors showed that FQs, when used systemically, are associated with disabling and potentially permanent serious side effects. These side effects can involve the disruption of tendons, joints, muscles, nerves, nervous system disturbances and even induction of type 2 diabetes. Due to the increasing number of reports about FQ toxicity and long-term complications, FDA has introduced significant restrictions on their use in recent years, particularly in children and in people aged 65 years.

1.1 Tendon rupture

FQs are associated with a significant risk of tendonitis and tendon rupture. Stephenson et al. [1] showed in their review the incidence of tendon injury among those taking FQs to be between 0.08 and 0.2%. In 2014, Lewis et al. [2] proved that FQ-related tendinopathy is a complication of treatment with this family of antibiotics and it is usually linked with 1 or more synergistic factors: male sex, age, renal disease, rheumatic disease, co-prescription of corticosteroid, and physical activity. For this reason, some sports medicine specialists have advised avoidance of FQs for athletes. Some authors e.g. [3, 4] proved that chronic renal disease, concomitant use of corticosteroids and age >60 years are known risk factors for FQ-induced tendinopathies. Concluding, FQs are associated with an increased risk of tendinitis and tendon rupture. This risk is further increased in those over age 60, in kidney, heart, and lung transplant recipients, and with use of concomitant steroid therapy.

1.2. Nervous system disturbances

Taking FQs is associated with their neurotoxicity, as well [5-8]. The main symptoms being correlated to FQ treatment include insomnia, restlessness, and rarely, seizure, convulsions, and psychosis [9-11]. Many reports point to chronic persistent peripheral neuropathy to be generated by FQs [12-18]. Cohen et al. [19] showed that a possible association between FQ and severe, long-term adverse effects involving the peripheral nervous system as well as other organ systems are observed.

1.3 Cardiotoxicity

Stahlmann et al. [20] showed that FQs prolong the heart's QT interval by blocking voltage-gated potassium channels. In some cases this can be a life threatening condition because of prolongation of the QT interval can lead to torsades de pointes, a life-threatening arrhythmia. Statistically significant risk factors for clinically significant changes in QTc were hypokalemia and a left ventricular ejection fraction < 55%.

1.4. Hepatotoxicity, nephrotoxicity

The other adverse reactions generated by FQs include hepatotoxicity [21], nephrotoxicity [22]. Golomb et al. [23] reported case-serious study and showed the potential occurrence of serious, persistent and delayed multi-symptome serious side-effects apparently triggered by FQ use causing severe functional compromise and disability in previously vigorous, healthy individuals. In this study, Golomb et al. described patients who developed new-onset symptoms during and following FQ use. Domains of serious and persistent sequels included the better recognized tendon and muscle issues, but extended to the well-reported but still often unappreciated potential for cognitive, psychiatric, peripheral nervous and gastrointestinal as well as endocrine issues.

1.5. Diabetes mellitus

Telfer et al. [24] conducted interesting study about FQs intake and development of type 2 diabetes mellitus (T2DM). They hypothesized that FQs induce an intracellular Mg²⁺ deficit that can lead to insulin resistance. Their data suggests that FQ exposure predisposes an individual to develop diabetes. He also showed a strong correlation between the increase in FQ application in U.S. in years 1990-2012 and the increase in T2DM incidences in subsequent years which suggests a large part of T2DM to be maybe generated by FQ exposure.

1.6. FQ-Associated Disability

In 2016 Kaur et al. [8] conducted basic science and clinical investigations of a newly identified adverse drug reaction, termed FQ-Associated Disability (FQAD). They proved that severe toxicities that develop when cancer patients receive supportive care drugs such as FQs are important, yet difficult to understand, detect, and to communicate to clinicians. Their findings supported recommendations of the FDA's advisory committee. Revision of FQ-product labels should be considered to include prominent descriptions of a newly identified FQ-associated long-term toxicity.

Concluding, patients with impairments of the CNS (e.g., epilepsy or arteriosclerosis), prolongation of the QT, elderly persons and individuals with concurrent use of glucocorticoids or chronic renal diseases should not be treated with FQs. FQs are contraindicated in children because they cause destruction of the immature joint cartilage in animals. The use in pediatrics is restricted to life-threatening infections.

2. Oxidative stress

One of the main effects generated by FQs in cells is connected with the oxidative stress (OS). Thus, a brief review about OS is presented below.

The main aspect of OS involves the overdosed leakage of electrons from the electron transport chain (ETC). In the normal, healthy state, the Krebs Cycle (KC) supplies hydrogen from glyco- and lipolysis in the form of NADH_2 and FADH_2 from KC to ETC. ETC separates hydrogen into protons and electrons. Protons are transported into mitochondrial intermembrane space (IMS) generating proton gradient and mitochondrial membrane potential $\Delta\Psi_m$ across the inner membrane and electrons are transported into oxygen. This complex process is a masterwork of the evolution because 4 electrons must enter the oxygen simultaneously in order to produce 2 water molecules. In the case of the block in the electron transport and the oxygen not to be fully reduced in one step, the reactive oxygen species (ROS) are created, which might be able to generate OS.

The total physiological electron leakage earlier estimated to be about 1% of electrons supplied to ETC [25, 26], currently has been estimated in favourable conditions to be 0.1-0.5% in rest [27-29] and 0.01-0.03% in the exercise [30, 31]. Assuming the oxygen consumption by human body in the rest to be about 500g (which corresponds the basal metabolic rate of 1920kcal/24h) the total electron current in all human mitochondria can be easily estimated to be about 70 Amper. The 0.1%-leakage denotes the Leakage Electron Current (LEC) to be about 0.07 Amper. In the case of physical or mental exercise both the metabolic rate and LEC increase several times, however, recent studies show that LEC does not depend so strictly on the metabolic rate [30]. Disturbing the precise mechanism of electron flow through ETC causes nearly always the increased leakage which can increase LEC even up to 10% (~7A in the rest). Majority of toxins joining ETC may do this leak. Because the increased OS is the side effect of the exercise state and OS is one of limits of the exercise, the increase in LEC during the rest may reduce the reserve for the OS increase during the exercise. The patient may feel tired even during small physical or mental exercise.

O_2^- is the first oxygen radical created by LEC. Joining the second electron causes the generation of H_2O_2 and joining the third electron to H_2O_2 creates a very dangerous hydroxyl radical OH^* . H_2O_2 is relatively stable and works as cellular sensor of the OS state and is involved in the cellular metabolism regulation [30, 32]. Brand et al. [30] has recently detected 11 sites in ETC where electrons leak from ETC generating OS. Brand and coworkers propose new postulate that the rate of electron leak does not depend on the electron flow rate but on the redox state (electron pressure) of the given site of leakage. Thus, ETC blockers increase the leak in sites before the block and decrease the leak behind the block. Majority of the sites generate the leak to the mitochondrial matrix and only two sites to both sides of inner mitochondrial membrane: the sites III_{Q_0} connected with complex III and G_Q connected with glycerol 3-phosphate dehydrogenase [30]. Brand et al. suggest also that the increase in O_2^- production from complex III that signals hypoxia to the HIF-1 α signaling system in cells is probably caused by indirectly changed metabolite concentrations that deliver more electrons to the ETC and lead to higher electron leak to oxygen. This is of high importance because as discussed later in this paper, HIF-1 α signaling is blocked in FQ patients disturbing strongly very important energy production regulatory pathway.

The last radical being created during the OS is OH^* . It is very dangerous because his lifetime is very short (10^{-9}s) and the cell does not possess any enzymes removing it. The overdosed production of OH^* is generated especially by Fenton reaction or in the ischemia state in which, because of lack of the oxygen, all the electrons create ROS. This massive ROS production induces the death of the ischemic cell.

The relative high physiological LEC forced the evolution to create the mechanisms carrying against free radicals. The first and, thus, most important barrier is the enzyme SOD2 (MnSOD, mitochondrial superoxide dismutase) which annihilates O_2^- - the first molecule of the O_2 -radical chain ($2\text{O}_2^- + 2\text{H}^+ \rightarrow \text{O}_2 + \text{H}_2\text{O}_2$). H_2O_2 is next removed by catalase or glutathione peroxydase. Meanwhile, it comes out of mitochondria and regulates e.g. redox dependent Kv1.5 channels in the cell membrane [32]. This mitochondria-ROS-Kv channel axis is now recognized as basis of an important O_2 -sensing mechanism in many tissues [33].

The simplified scheme of the electron leakage from ETC is presented in Figure 1.




Figure 1. The schematic presentation of the ATP production system. The acetyl-CoA supports Krebs Cycle to produce NADH_2 . Hydrogen from NADH_2 (and FADH_2) enters the cytochrome chain. Some electrons leak before they reach oxygen generating O_2^- and next H_2O_2 . H_2O_2 comes away from mitochondria and works as redox signaling molecule. The amount of leaking electrons depends mainly on NADH_2/NAD ratio (hydrogen pressure). The degree of PTP opening must be precisely regulated and depends on many factors, e.g. $\text{O}_2^-/\text{H}_2\text{O}_2$, Ca^{2+} , $\Delta\Psi_m$.

2.1. The role of Mitochondrial Permeability Transition Pores (PTP) in the regulation of energy production.

The mitochondrial Permeability Transition Pore (PTP) is the large protein complex placed in the outer mitochondrial membrane being precisely regulated by many factors [34, 35]. It consists mainly of voltage-dependent anion channel (VDAC), adenine nucleotide translocase (ANT) and cyclophillin D (CypD). In order to make the cell function properly, the degree of opening state of this complex must be precisely fitted to the actual physiological state of the cell. If the complex is open, the non-selective traffic between the IMS and cytosol of small-charged particles, water and substances up to 1.5 kDa takes place. ADP can enter mitochondria to produce ATP but the protons leak from IMS to cytoplasm reducing the mitochondrial potential $\Delta\Psi_m$ from -140mV to about -110mV and contributing to apoptosis. If the complex is closed, ADP cannot enter mitochondria to produce ATP and $\Delta\Psi_m$ grows from -140 to -160mV. The decrease in $\Delta\Psi_m$ is characteristic for OS state and the increase in $\Delta\Psi_m$ is characteristic for some types of cancers [32]. This observation explains the ability of FQs to treat cancers [36-41].

Main factors that regulate PTP state [34]:

- Opening PTP:
 - $[Ca^{2+}]_{mit}$: concentration of mitochondrial Ca^{2+}
 - Reduced $\Delta\Psi_m$ (positive loop!)
 - Free radicals (oxidative stress)
 - Inorganic phosphate (only with Ca^{2+})
 - Some apoptotic factors
- Closing PTP:
 - Acidic pH (a part of cancer state is lactate accumulation in the cell)
 - ATP, ADP, NADH
 - Mg^{2+}

Hexokinase II (HKII), mitochondrial creatine kinase (CK), benzodiazepine receptor (PBR), and Bcl-2-family members (Bcl-2, Bcl-xL, and Bax) are putative regulatory components.

It can be observed that OS contributes to opening PTP and reducing $\Delta\Psi_m$. Reduced $\Delta\Psi_m$ causes further opening of PTP and the decrease in energy production. The final step and physiological sense of this positive loop is the induction of the apoptosis. However, if the apoptosis induction is not reached, the new balance between factors being in the common regulatory loop finds new equilibrium point which can be far away from that optimal one. The 'positive loop' regulation of PTP points to its high sensitivity to different factors controlling it and suggests the regulation of PTP to be one of the most important points of energy production control. Some positive regulatory loops are presented in Figure 2. According to the Control Theory, such positive loops denote that the reaction to small stimulus may be strongly magnified. The influence of FQs on the detailed regulation of PTP is the urgent topic for further research.

2.2. How the cell adapts to OS state?

Increased OS state is characterized first by increased H_2O_2 level in the cell. In the physiological state, H_2O_2 level informs the cell and nucleus about the mitochondrial energy production state because, in the physiological conditions, it is connected with the metabolic rate of the cell. In the case of disturbed metabolic regulation (e.g. mitochondrial toxins), increased OS and open PTP - ATP production is reduced. The main process that consumes ATP in the cell is the Na/K pump that removes Na^+ from the cell, pumps K^+ to the cell and generates the negative membrane potential of the cell. The work of Na/K pump is the most fundamental process of life because it consumes 20% to 50% of ATP and generates strongly negative charge inside the cell. Many other transports that take place across the cell membrane work as Na^+ -co-transport, e.g. amino acids, phosphates, Ca^{2+} , boric acid. One molecule of Na^+ co-transport contributes to about 30% of ATP/ADP Gibbs free energy, thus, it can be treated as 'small energy bricks' when compared to ATP as a large one. The lack of ATP reduces the work of Na/K-pump reducing negative membrane potential ΔV and K^+ concentration in the cell. However, H_2O_2 generated during OS activates the opening of redox dependent Kv1.5 channels in the cell membrane [32] contributing to the further efflux of K^+ from the cell. The escaping K^+ increases temporarily the negative potential inside the cell. It can be concluded that restoring the negative cell potential is more important for the cell than keeping high K^+ concentration in the cell.

On the other hand, in the case of reduced ATP production, the lowered ΔV opens voltage-gated Ca^{2+} channels which causes Ca^{2+} influx into the cell [32]. Increased $[\text{Ca}^{2+}]_i$ activates calcineurin. Activated calcineurin shifts NFAT (Nuclear Factor of Activated T cells) to the nucleus where it inhibits Kv1.5 potassium channels production. Lowered amount of Kv1.5 channels reduces K^+ efflux. Concluding, in the OS state Kv1.5 channels are more open, however, their amount is reduced. Two opposite mechanisms regulating the membrane potential and K^+ efflux find some new equilibrium. This mechanism seems to be the natural adaptation process to the increased metabolic rate of the cell generating physiological OS. In the state of increased metabolic rate the concentration of K^+ (and most probably Mg^{2+}) is reduced, the concentration of Ca^{2+} and H_2O_2 (and most probably Na^+) is higher, the membrane potential is also reduced. If the healthy cell returns from the increased metabolic rate to the resting state all these parameters return to their optimal values. This return depends on the ability of the mitochondria to produce ATP which drives Na/K-pump.

In the case of the permanent OS or disturbed state of PTP being inadequately open, this return cannot take place because the ATP production is lowered. The cell being in the resting state, comes to the state which can be called "permanent stress adaptation". In the case of the necessity to increase the metabolic rate, the further increase in metabolic rate is difficult because of the lack of physiological adaptation reserve. The final effect for the patient is the feeling of "the lack of energy". Many other regulative processes take place, of course, as OS adaptation. However, the above described changes belong in authors opinion, to the primary regulatory axes.

3. Molecular mechanisms of FQ toxicity

Good understanding of the OS state is very important for understanding the consequences of OS generation by FQs. From the therapeutic point of view, the important question concerns the molecular mechanisms leading FQs to generate OS, because they determine the possible effective treatment of this state. Many people are waiting for the understanding of the FQ-toxicity mechanisms and treatment methods. Beside OS, epigenetic effects of FQs are of high importance, as well. The epigenetic effects may depend on the methylation of DNA and/or histones, however, ROS contribute also to epigenetic changes [42]. Some authors point also to the similarity of bacterial and mitochondrial DNA, both existing in circular super-twisted helices and gyrase-like enzymes being postulated to be responsible for the organization of mitochondrial DNA, suggesting the possible direct effect of FQs to mitochondrial DNA leading to the disturbed mitochondria regeneration and division [43, 44]. The changes in the cytoskeleton were observed also after FQ treatment [45] and cytoskeleton has been demonstrated to be strictly connected with energy dissipation and organization in mitochondria [46-49]. The most important elements of FQ toxicity are presented in Figure 2. The positive regulatory loops magnifying the toxic effects are marked with '+'. Let us analyze the most important aspects of the molecular activity of FQs in the cell, being reported until now.




Figure 2. The main ways of FQ toxicity. The positive regulatory loops magnifying the toxicity of FQs are marked with '+'. The '?' signs denote the possible but not confirmed effects of FQ toxicity.

3.1. Chelating bivalent cations and proteins by FQs

FQs are the group of chemical compounds called zwitterions. Zwitterion is a neutral molecule with both positive and negative electrical charges on its opposite sides. This feature makes them possible to create strong complexes with both protein anions and positively charged bivalent cations. The protonation constants for the acidic and base part of the FQ molecules were estimated as $pK_1=8.2-8.5$ and $pK_2=5.6-6.2$. These values denote that the dissociation coefficients equal about 90-95% in neutral intracellular conditions of $pH=7.0$ for both the acidic and base side of the molecule [50]. The structure of the exemplary FQ is presented in Figure 3.




Figure 3. The exemplary FQ - norfloxacin and its zwitterion structure.

FQs possess two main sites for metal chelate formation. The first one, represented by the carbonyl and carboxyl groups in neighboring positions, is the most common coordination mode in the quinolone chelates [50]. Quinolones bind divalent cations as Mg^{2+} , Ca^{2+} , Cu^{2+} , Zn^{2+} , Fe^{2+} , Co^{2+} , forming chelates with 1:1 or 1:2 metal:FQ stoichiometry or trivalent cations (e.g. Al^{3+} , Fe^{3+}), forming chelates with 1:1, 1:2 or 1:3 metal:FQ stoichiometry. The constant values for CIP chelates have been measured to decrease in the order: $Al^{3+} > Fe^{3+} > Cu^{2+} > Zn^{2+} > Mn^{2+} > Mg^{2+}$ [51]. For NOR chelates, the relation is quite similar: $Fe^{3+} > Al^{3+} > Cu^{2+} > Fe^{2+} > Zn^{2+} > Mg^{2+} > Ca^{2+}$. Let us observe that presented research [50, 51] did not analyze Se^{2+} ions which are also of high importance in the cell, because Se^{2+} is the co-factor of glutathione peroxidase removing H_2O_2 . The examination of the ability of FQs to generate Se^{2+} -FQ complexes is the important aim for further research.

Seedher et al. [52] analyzed the ability of 5 cations: Fe^{3+} , Al^{3+} , Zn^{2+} , Cu^{2+} and Mg^{2+} to create the complexes with four FQs and human serum proteins using fluorescence UV absorption spectroscopy. He measured the association constants to be of the order of 10^2-10^4 for the FQ- Me^{n+} interaction. The interaction was the highest for Al^{3+} and lowest for Mg^{2+} . At a Me^{n+} /drug ratio of 1:1, approximately 50%-73% of metal ion was bound per mole drug in the most cases. Seedher's results indicate that chelate formation with bivalent metals can cause significant alteration in the human serum-FQ binding affinity.

Koga et al. [53] measured the FQ concentration ratio between intra- and extracellular fluid in human polymorphonuclear leukocytes using high-performance liquid chromatography. This c_i/c_{out} ratio oscillated between 2.2 to 8.2. In another paper Pascual et al. [54] presented c_i/c_{out} gradient for trovafloxacin (TRV) to be about 9 for extracellular concentrations ranging from 0.5 to 25 ug/ml. Assuming net charge of the FQ molecules being close to 0, the expected ratio for the soluble fraction is expected to be close to 1:1. These results point to the ability of FQs to create complexes with different intracellular molecules which do not participate to the soluble fraction of FQs in the cell.

The other important feature of FQs has been presented by V.T. Andriole et al. [55]. Namely, they estimated the minimum solubility of FQs in neutral pH. They pointed that this class of molecules is characterized by very high melting point, generally $>200^{\circ}\text{C}$, which indicates that the crystal forms are very stable.

All these FQ features strongly support the thesis that FQs can survive in the cell for a long time contributing to chronic, long-term adverse reaction in patients treated with FQs. The question, to what extent takes this phenomenon place and if it contributes to chronic symptoms of FQAD, remains unclear.

Metal ions chelating seems to be the most fundamental feature of FQs which probably leads to all other observed toxic effects. The antibacterial effect is connected with chelating Mg^{2+} which disturbs the gyrase and topoisomerase interaction with DNA. However, the Mg^{2+} is described to create weaker chelates with FQs than other important ions like $\text{Fe}^{2/3+}$, $\text{Cu}^{1/2+}$, Zn^{2+} and Mn^{2+} .

The well described activity of FQs which concerns the Fe^{3+} chelating was examined by Badal et al. [56]. They examined that three FQs: NOR, CIP and enrofloxacin (ENR) are the powerful $\text{Fe}^{2/3+}$ chelators comparable with deferoxamine, a clinically useful Fe-chelating agent. He showed that $\text{Fe}^{2/3+}$ chelating by FQs leads to epigenetic effects through inhibition of α -dependent dioxygenases (DOXG) that require Fe as co-factor. Three important DOXGs were analyzed: jumonji domain histone demethylase, TET DNA demethylase and collagen prolyl 4-hydroxylase. The activity of all 3 enzymes was reduced by FQs in micromolar concentrations leading to accumulation of methylated histones and DNA and to inhibition of proline hydroxylation. The IC_{50} concentrations for Fe-chelation was equal to about $52 \pm 20 \mu\text{M}$ for CIP, $44 \pm 15 \mu\text{M}$ for NOR, $41 \pm 20 \mu\text{M}$ for ENR and $360 \pm 25 \mu\text{M}$ for deferoxamine. These results point to high ability of FQs to absorb $\text{Fe}^{2/3+}$ and reduce the activity of enzymes which use $\text{Fe}^{2/3+}$ as co-factor. The results suggest also the possibility that the tendon FQ-toxicity is dependent on lack of collagen proline hydroxylation which changes significantly the mechanical properties of the collagen. Some other studies report, however, the association of the enhanced extracellular matrix metalloproteinases [51, 52], mainly collagenases expression [53], to be associated with FQ-induced tendinopathy.

It must be pointed that cytochromes are the other important proteins which use $\text{Fe}^{2/3+}$ in hem groups as a co-factor. Thus, the question arises if the reduced free $\text{Fe}^{2/3+}$ concentration in the cell contributes to ETC inhibition, electron leakage and/or OS? The question arises, as well, which cellular effects are connected with Cu^{2+} and Zn^{2+} ions, because Cu^{+} is also an important cytochrome co-factor and Zn^{2+} is important due to high number of enzymes (about 300 total)

being co-factored by this element and the total cytoplasm concentration of Zn^{2+} is similar and even higher than that of $Fe^{2/3+}$. Valko et al. [57] presents that redox-inert zinc (Zn^{2+}) is the most abundant metal in the brain and an essential component of numerous proteins involved in biological defense mechanisms against oxidative stress. The depletion of zinc may enhance DNA damage by impairing DNA repair mechanisms.

Some effects of Mg^{2+} -chelation are also well described, especially with respect to the cartilage damage. Mg^{2+} is an important intracellular ion being a co-factor of about 300 enzymes. Shakibaei et al. [58] showed that effects of individual dose of ofloxacin gives identical effects in cartilage observed in electron microscopy as Mg^{2+} deficient diet suggesting that quinolone-induced arthropathy is probably caused by a reduction of functionally available Mg^{2+} in cartilage. Similar results were observed during cultivation of chondrocytes in Mg-free medium [45, 59-61]. The supplementation of Mg^{2+} accompanying the FQ treatment restored to some degree the cartilage lesions [60, 61], however, did not restore the reduced cell division [61]. This suggests that other mechanisms are involved in cell division reduction after FQs [44]. Mg^{2+} deficiency in immature dogs induced similar clinical symptoms as quinolone treatment: distinct alterations in chondrocytic fibronectin staining and their ultrastructure [62, 63]. The effect of Mg^{2+} supplementation had a two-way effect on FQ-treated cultured chondrocytes: the number of cells adhering to culture support was increased and cell morphology was comparable to that of control cells. This suggests that addition of Mg^{2+} restores extracellular Mg^{2+} -dependent inter-cell interactions.

On the other hand, dietary Mg^{2+} is also presented to reduce intestinal FQ absorption [64-70] and it is also postulated to be in relation to diabetes type 2 and FQ treatment [24].

Summing up, the number of enzymes possessing reduced activity due to their ion-cofactor chelation is probably long and it is the important topic for further research. The separate problem consists the chronicity of ion-chelation by FQs. The presented research does not describe the chronic state of FQAD but the phenomena taking place during FQ application. It must be analyzed, to which degree persistent ion chelation takes place at FQAD patients.

3.2. Oxidative stress generated by FQs

Many papers point to the feature of FQs to generate oxidative stress in the cells. The molecular mechanisms leading to OS state differ probably in details for different FQs depending on different ability to chelate subsequent metal ions and on possible different ability to change enzymes activity in the ion-independent manner.

An example of ion-independent FQ activity was presented by Qin et al. [71]. He analyzed the influence of CIP and ENR on the erythrocytic catalase (CAT), a vital enzyme involved in OS reduction (CAT reduces H_2O_2 to O_2 and H_2O). The cellular tests firstly confirmed an enhanced oxidative stress in FQ-treated erythrocytes in the form of the GSH contents depletion and decrease in CAT activity with CIP effect to be more harmful than ENR. Next, spectroscopic computations showed the FQ binding places to CAT takes place mainly through electrostatic forces. Binding of two FQs not only caused the conformational and micro-environmental changes of CAT, but also inhibited its molecular activity, which was consistent with the cellular activity measurements. On the other hand, the treatment with danlofloxacin

[72] increased antioxidant enzymes activities such as superoxide dismutase (SOD) and catalase (CAT), suggesting that the ability of subsequent FQs to change the activity of different antioxidative enzymes can vary significantly.

Many papers present the existence of OS induced by FQs, e.g. Pouzard et al. [73] measured the redox status change of immortalized rabbit tendon cells as a response to pefloxacin, ofloxacin, LEV and CIP. All FQs showed moderate cytotoxicity on tendon cells after 24h and more severe, significant toxicity after 72h. The intracellular redox potential has been reduced slightly but significantly after 72h even at concentrations 1 μ M (~10 μ M is the therapeutic one) and strongly reduced at concentrations 1mM (100x higher than therapeutic ones). ROS production has increased slightly (~25%) but significantly at therapeutic conditions and strongly (~150%) at 100 μ M (10x higher than therapeutic ones). The intracellular GSH concentration was reduced by 20-50% even at 0,01 μ M concentrations (1000x smaller than therapeutic ones) and the collapse (decrease in 50-90%) in GSH was observed at 1mM. The question arises if the GSH depletion is connected only with the increased H₂O₂ generation or also with a reduced activity of GSH-reductase which restores the reduced form of GSH (GSSG+NADPH₂→2GSH+NADP).

It is important to observe that the rapid increase in the toxicity takes place after the given concentration of FQs is reached which is only slightly higher than the therapeutic one. The substantial increase in the ROS production which can lead to serious consequences begins at concentrations being approximately 10x higher than that therapeutic ones. This factor is, however, estimated with low precision because no intermediate concentrations were examined between 0.01, 0.1, 1, 10, 100 and 1000 μ M. Assuming that some people are charged with reduced ROS annihilation ability (e.g. as presented in [21]), this toxicity limit may occur at lower, therapeutic concentrations. This experiment stays also in the agreement with clinical observations that the FQAD takes place especially in patients who were treated with higher FQ doses, by a longer period or with FQ series repeated a few times in a short period of time. In the other paper, Pouzard et al. [74] observed in vitro in rat tendons the increased ROS production and reduced GSH concentration. Similar results were also showed by Yu et al. [72].

Some papers point to detailed FQ effects on different enzymes. Hsiao et al. [21] found that TRV induced OS on heterozygous SOD2(+/-) deficiency mice was higher than on the normal mice. Hepatic protein carbonyls were increased by 2.5-fold and hepatic mitochondrial aconitase activity was decreased by 20% in mutant, but not in wild-type mice. Because aconitase is a major target of peroxynitrite, they determined the extent of nitrotyrosine residues in hepatic mitochondrial proteins. (Peroxynitrite ONOO⁻ is formed by the combination of O₂⁻ and NO.) TRV significantly increased nitrotyrosine in SOD2(+/-) mice only. TRV increased also the production of mitochondrial NO in immortalized human hepatocytes. Similarly, mitochondrial Ca²⁺ was increased by TRV, suggesting Ca²⁺-dependent activation of mitochondrial NOS activity. Furthermore, the transcript levels of the mtDNA-encoded gene Cox2/mtCo2 were decreased in SOD2(+/-) mice only, while the expression of nDNA-encoded mitochondrial genes was not significantly altered in both genotypes, suggesting selective effects on mtDNA expression. These data indicate that TRV enhances hepatic mitochondrial peroxynitrite stress at increased basal O₂⁻ levels, leading to the disruption of critical mitochondrial enzymes and gene regulation.

Another important information can also be found in [21]. OS generation by FQs may take place at lower FQ concentrations in some people being charged with lower ability to reduce OS. The reasons of reduced OS barrier may be different, however, the most important seem to be the heterozygous mutations, trace element deficiencies and charge of the cells with other toxins contributing to OS.

Kumbhar et al. [75] reported gatifloxacin (GAT) to produce retinal damage in rabbits and significant alteration in the antioxidant status indicated by the decreased activity of superoxide dismutase and decreased levels of blood glutathione with a concomitant increase in the activity of catalase, glutathione peroxidase, and glutathione S-transferase enzymes. The levels of malondialdehyde were also elevated. The effects were dose-dependent. Talla et al. [76] examined the OS parameters in the blood after CIP, LEV and GAT therapy on SOD3 (extracellular), glutathione, plasma antioxidant status and lipid peroxides evaluated at 53 patients on different dosage regimens up to 5 days. The significant elevation of lipid peroxide was observed in patients treated with CIP and LEV. The substantial depletion in SOD3 and glutathione was observed especially in CIP patients. All three FQs reduced the plasma antioxidant status, but especially CIP and LEV.

Liu et al. [77] determined the effect of ENR on the release of lactate dehydrogenase (LDH), reactive oxygen species (ROS), superoxide dismutase (SOD), total antioxidant capacity (T-AOC), malondialdehyde (MDA), mitochondria membrane potential $\Delta\Psi_m$ and apoptosis in the hepatic cell line of grass carp. The doses of 50, 100 and 200 $\mu\text{g/mL}$ increased the LDH release and MDA concentration, induced cell apoptosis and reduced the $\Delta\Psi_m$ compared to the control. The highest dose of 200 $\mu\text{g/mL}$ also significantly reduced T-AOC.

All the presented above experiments show the increased OS state after FQ treatment. The changes in enzymes activity varies between experiments, tissues and kinds of FQs suggesting possible variability of the common relations. What seems to be important is the reduction in SOD activity which is the first line barrier against O_2^- . New experiments must estimate these changes in details, however, more attention must be paid to mitochondrial MnSOD which cares against mtDNA damage by O_2^- generated by leaking electrons from ETC to mitochondrial matrix. The increased activities of some anti-OS enzymes seem to be the OS adaptation processes.

3.3. Reduction of mitochondrial $\Delta\Psi_m$ potential by FQs

One of the cellular symptoms present in the FQ-charged cells is the reduction of the mitochondrial potential $\Delta\Psi_m$ [37, 38, 77-79], however, the detailed mechanism of this phenomenon remains unknown. Since the main mitochondria uncoupling factor is the PTP, the reasons of the reduced $\Delta\Psi_m$ should be searched between factors that regulate PTP opening. The first possibility is the OS by itself being generated by FQs. ROS can induce VDAC oligomerization (main part of PTP) to yield a mega-channel creation which are postulated to create large holes being able to release cytochrome c to cytoplasm. O_2^- -induced apoptosis has been found to be inhibited by DIDS or anti-VDAC antibodies (VDAC blockers), which suggests that O_2^- increases VDAC-dependent permeabilisation of the outer mitochondrial membrane [80, 81].

One of the proteins which can support PTP opening is Translator Protein (TSPO), called also peripheral-type benzodiazepine receptor or isoquinoline binding protein. TSPO is predominantly located on the surface of the mitochondria where it is postulated to physically associate with VDAC-ANT. It has been suggested that TSPO may activate PTP opening, causing $\Delta\Psi_m$ reduction and leading to apoptosis [80, 81].



Some authors suggest that epileptogenic activity of FQs possibly relates to GABA-like structure of some FQs which may allow them to act as GABA antagonists [82, 83]. Since TSPO is also a benzodiazepine receptor, similar interaction may maybe also take place between FQs and TSPO leading to opening PTP.

The problem of the PTP opening is probably, however, more complicated and it is in relation with many other factors involved in energy production and apoptosis induction in the cell. The broad reviews about regulation of PTP and VDAC (the main PTP protein) are presented in [35] and [34].

The other way to reduce $\Delta\Psi_m$ is the opening of UCP channels which generate the proton leak through the inner mitochondrial membrane. UCP regulation is a separate problem and possible way for treating FQAD patients. The review of this problem is published by Divacaruni et al. [84].

3.3. Does Fenton reaction take place in FQAD patients?

The next problem that is connected especially with Fe^{2+} and Cu^+ ions is the possibility of the Fenton reaction (FR) to be generated on $\text{Fe}^{2/3+}$ and $\text{Cu}^{1/2+}$ ions, but maybe on the FQ- Fe^{2+} complexes, as well. Fenton reaction consists in the conversion of O_2^- and H_2O_2 into strongly dangerous OH^* radical, it takes place on $\text{Fe}^{2+/3+}$ and $\text{Cu}^{1+/2+}$ ions and consists of 2 steps:



This reaction increases strongly the effects of OS in the cell and if it is too intense, it may contribute to cell death. The question arises: to what degree does the Fenton reaction magnify initial OS effects?

There is no evidence that would prove such reaction to take place in FQAD patients, however, four theses could be postulated for further research: (a) The reaction is increased just due to the increased substrates concentration (O_2^- and H_2O_2) for FR. (b) The reaction intensity is reduced with respect to that expected due to the reduced Fe^{2+} level in the cell. This can take place due to reduced membrane potential and reduced ability Fe^{2+} to be pulled into the cell. (c) The reaction is magnified due to the increased ability of FQ- Fe^{2+} complexes to generate OH^* in FR reaction by itself. (d) The reaction is magnified due to up-regulation of Fe-transporting protein which increases Fe^{2+} concentration in the cell. Similar upregulation has been detected in *Streptococcus pneumoniae* by Ferrandiz et al. [85, 86]. They observed the

upregulation of the genes of the fatDCEB operon, involved in iron (Fe^{2+} and Fe^{3+}) uptake. In accordance, they observed an attenuation of LEV lethality in iron-deficient media. However, the bacterial gene regulation cannot be directly compared to that mammalian one. On the other hand, electro-Fenton reaction is described to perform degradation of LEV in experimental conditions [87, 88], however, it seems to be of low probability, such reaction to take place at in-vivo conditions.

3.4. Changes in gene expression and enzyme activities after FQ treatment

Besides OS aspects connected with FQAD, some papers point to other effects of FQ toxicity. Fox et al. [89] measured reverse-transcriptase quantitative polymerase chain reaction analyses on total RNA isolated from supraspinatus tendon of rats. He showed the significant upregulation of IL-1b mRNA, Tumor necrosis factor (TNF), Matrix metalloproteinases MMP-3 (30x increase), MMP-13 (7x) and the tissue inhibitor of metalloproteinases (TIMP)-1 (4x) in the FQ treated rats. FQ-treated groups showed significantly less fibrocartilage and poorly organized collagen at the healing enthesis compared with control animals.

Aranha et al. [41] measured the gene expression effects of CIP on prostate carcinoma and healthy control cells. Treatment of prostate cancer cells with CIP resulted in a dose- and time-dependent inhibition of cell growth (70-100% with 50-400 ug/ml). Cells were arrested at the S and G2/M phases and apoptosis was induced. The cyclin-dependent kinase (CDK) inhibitor p21/WAF1 was down-regulated 12h following CIP in treatment which can lead to rapid CDK2 activation and caspase induced apoptosis. There was also observed a significant increase in the Bax/Bcl-2 ratio with translocation of pro-apoptotic Bax to mitochondria and activation of caspase-3. Let us recall that Bax, Bcl-2 and caspase-3 are involved in PTP opening state.

Badal et al. [56] showed that NOR, CIP, and ENR as iron chelators leads to epigenetic effects through inhibition of alpha-ketoglutarate-dependent dioxygenases that require iron as a co-factor. Three dioxygenases were examined in HEK293 cells treated with FQ. At micromolar concentrations, these antibiotics inhibited jumonji domain histone demethylases, TET DNA demethylases, and collagen prolyl 4-hydroxylases, leading to the accumulation of methylated histones and DNA and inhibition of proline hydroxylation in collagen, respectively. These effects may explain FQ-induced nephrotoxicity and tendinopathy. Let us observe that the changes in DNA and histone methylation state cause strong and broad epigenetic effects being difficult to predict.

Liang et al. [90] measured the sub-chronic toxic effects of NOR on a swordtail fish by measuring mRNA expression of cytochrome P450 1A (CYP1A), cytochrome P450 3A (CYP3A), glutathione S-transferase (GST) and P-glycoprotein (P-gp) and their corresponding enzyme activities. Results showed that NOR significantly affected the expression of CYP1A, CYP3A, GST and P-gp genes in swordtails. The gene expressions were, however, more responsive to NOR exposure than their corresponding enzyme activities. The analyzed enzymes are very important because they express the ability to catalyze detoxification of xenobiotic substrates, including FQs. The possible reduction of its activity in humans may be of high importance, because FQs undergo biotransformation in the liver from approximately

50 percent for pefloxacin to about 6 percent for ofloxacin [91]. Although glucuronide conjugates have been identified as minor metabolites for some agents, most metabolic reactions involving quinolones occur through microsomal oxidative mechanisms at the cytochrome P-450 site. These metabolic alterations involve the piperazinyl moiety and usually result in compounds with significantly less microbiologic activity than the parent drugs. However, the conclusions from fish cannot be directly transferred to humans, the results suggest the possibility of the delayed toxicity to be connected with reduced detoxification induced by itself.

Similar results of P-450 inhibition by FQs were found by Shlosberg in chicken [92] and Granfors [93]. Regmi et al. pointed to the inhibiting effect of FQs in dogs on P-450 1A but not on P-450 3A [94, 95].

3.5. FQs and HIF-1 α

Dioxygenase inhibition by FQs was predicted by Badal et al. [56] to stabilize transcription factor HIF-1 α by inhibition of the oxygen-dependent hypoxia-inducible transcription factor prolyl hydroxylation. In dramatic contrast to this prediction, HIF-1 α protein was eliminated by FQ treatment. Badalet coworkers explored this effect to be caused by inhibition of HIF-1 α mRNA translation.

The natural function of HIF-1 α system is to change the metabolism of the cell into the anaerobic pathway in order to protect the cell against OS. The conversion of pyruvate to lactate is enhanced by up-regulation of the lactate dehydrogenase-A (LDH-A). On the other hand, pyruvate dehydrogenase complex is inhibited by HIF-1 α by upregulating pyruvate dehydrogenase kinase (PDK) which inhibits PDH-A. Thus, HIF-1 α inhibits pyruvate to enter KC and to produce NADH₂. As presented by Brand [96], NADH₂/NAD ratio (which can be called 'hydrogen pressure') is the main factor determining the escape of electrons from the ETC chain, making this process to high degree independent of the total ETC electron flow. (The term 'hydrogen pressure' reflects the natural similarity of ETC to the water flowing through the pipe with small holes. The water escaping through the holes depends on the pressure but not on the water velocity in the pipe.) Thus, in case of the lack of the oxygen in the tissue which is the final recipient of the electrons from NADH₂, HIF-1 α turns on the side way for hydrogen, not to put it into ETC and produce OS but to produce lactate. The lack of this safety valve (HIF-1 α) in FQ patients may cause the shift of the overdosed amounts of hydrogen into ETC causing overdosed electron leak. Assuming this phenomenon to be important in FQ-treated patients, the glyco- and lipolysis inhibitors could be considered as OS reductors. The natural glycolysis inhibitor in the diet is citric acid which inhibits phosphofructokinase (main regulatory point of glycolysis) and activates fructose 1,6 biphosphatase which catalyses the opposite reaction promoting gluconeogenesis and pentose cycle pathway. On the opposite, the vinegar (acetic acid) could be contradicted by FQAD patients because it promotes glyco- and lipolysis both contributing to possible overdosed 'hydrogen pressure' and OS.

4. Therapeutic conclusions

The treatment of the FQAD, especially that lasting for many years, is a very difficult therapeutic problem. The effectiveness of different therapies carried out on patients is rather low. A large number of patients suffers from chronic tiredness, tendinopathies, neuropathies, need more sleep, even more than 12h/24h. Understanding all the molecular mechanisms of the FQ activity in the cell is the urgent aim for current science to find methods helping these people.

The main question that arises here, concerns the reasons of chronicity of FQAD symptoms which last for many years, sometimes, even after a standard 5-days FQ treatment. 3 reasons can be taken into consideration:

A. Long lasting OS destroys the mitochondrial DNA and the newly synthesized proteins creating cytochrome complexes are disturbed in their structure leading to permanent electron leakage and OS.

B. The complexes of FQ with proteins and cations are so stable that they exist in the cells by many years disturbing energy production and epigenetics.

C. Epigenetic changes in gene regulation become persistent many years of FQ application even in the case of lack of FQ in the cell.

The answer, which of these three possible reasons contribute to the chronicity of FQAD symptoms, is of high importance with respect to the problem of effective treatment of this state. The research answering these questions must be performed as fast as possible.

In the case of mtDNA destroying, the treatment is difficult and it must focus on the stimulation of mitochondrial replication. The more destroyed mitochondria must be removed and the less destroyed must replicate in order to substitute for the removed ones and to reduce the total LEC. After many replications, the most healthy mitochondria would dominate the cell. The final effect would depend on the state of the most healthy mitochondrion in the cell. The second possibility is to increase the ratio of cell exchange in the given tissue. The cells with more destroyed mitochondria must be shifted to apoptosis while more healthy cells must substitute them. This process, however, cannot take place in the Central Nervous System and muscles because the cell exchange is close to zero in these tissues. Also, the collagen exchange is very low causing the tendon regeneration to be difficult and long-lasting problem.

If new research would confirm the existence of FQs in the cells and mitochondria in the amounts making possible their permanent interactions with proteins and cations even after many years of FQ application, the research must focus on methods, how to remove FQs from strong protein- and cation complexes. The simplest way seems to be the application of increased doses of metal cations Fe^{2+} , Cu^+ , Mn^{2+} , Zn^{2+} , Mg^{2+} which are natural FQ-competitors for protein binding sites. It should be pointed that bivalent metal ions enter the cell to some degree due to the negative membrane potential of the cell and next, enter the mitochondria due to the negative value of mitochondrial potential $\Delta\Psi_m$. The Nernst equation $\Delta V = RT/zF \cdot \ln(c_{in}/c_{out})$ defines the equilibrium between the ion concentration gradient and voltage across the membrane. If the membrane voltage $\Delta V/\Delta\Psi_m$ decreases, the concentration

equilibrium gradient c_{in}/c_{out} of bivalent ions decreases significantly as well, because the lowered potential may not be able to pull bivalent cations into the cell/mitochondrium up to the required concentration. Table 1 shows the exemplary calculations.

$\Delta V/ \Delta\Psi_m$	Equilibrium c_{in}/c_{out} gradient ($z=2$ and $T=310K$)
-160mV	160.000x
-140mV	36.000x
-110mV	3.800x
-90mV	850x
-70mV	190x
-50mV	42x
-30mV	9x

Table 1. The conversion of Nernst equation $c_{in}/c_{out} = \exp(\Delta V \cdot zF/RT)$ for bivalent ions as Mg^{2+} , Mn^{2+} , Fe^{2+} , Zn^{2+} shows strong relation between the actual membrane potential and ability of the cell/mitochondrium to attract and absorb the ions into the cell/mitochondrium. Reduced membrane potential ΔV is a strong factor which hinders entering bivalent cations to the cell. However, the detailed mechanisms of individual ion transport must be analyzed.

The Table 1 shows that the decrease in membrane voltage reduces strongly the ability of the cell to pull X^{2+} ions into the cell/mitochondrium. The question arises, to what extent the transport of X^{2+} ions into the cell is disturbed in FQAD patients? And next, if the possible lower X^{2+} concentration depends only on reduced $\Delta V/\Delta\Psi_m$ or maybe also on the disturbed membrane transport being blocked by FQs joining to metal binding sites of transport proteins? Every analyzed bivalent cation requires a separate analysis.

The third possible reason of the permanent FQAD state is the permanent disorder in gene expression caused by some positive loop regulations. E.g. reduced Fe^{2+} level disturbs oxoglutarate dependent dioxygenases and increases methylation of DNA and histones which leads to reduced Fe^{2+} absorption to the cell. Many other loops are possible which could cause the chronic state of the patient despite the lack of FQs in the cell. They have to be recognized in order to find the methods restoring the normal regulatory state. This case is the most hopeful for patients, because e.g. the mtDNA damage is rather difficult to be effectively treated and gene expression regulation is difficult, however, possible.

Until detailed knowledge concerning FQ toxicity would be recognized, the following directions in supporting FQAD patients are proposed according to the known and probable mechanisms of FQ toxicity:

- A. reduction of the oxidative stress;
- B. restoring reduced mitochondrial potential $\Delta\Psi_m$;
- C. supplementation of uni- and bivalent cations that are chelated by FQs;

- D. supporting the mitochondrial replication in the cell - pulling the more destroyed to apoptosis and proliferation of the more healthy ones;
- E. removing FQs permanently accumulated in the cells (if this phenomenon takes place);
- F. regulating the disturbed epigenetics and enzyme activities.

A. Assuming that H_2O_2 is not effectively removed from the cell after FQ treatment, subsequent consequences may occur such as opening Kv1.5 channels, Fenton reaction, peroxynitrite radical creation, opening PTP channels, decoupling mitochondrial potential and finally, breaking down OS barrier. The detailed comparison of different FQs with respect to OS is urgent to determine which FQs are safer in use and which ones are more dangerous. Reduction of OS is a very broad area. There are thousands of natural substances which possess the antioxidant capacity and which are able to reduce free radicals leaked from ETC. One should, however, remember that they work as one to one. It means that, as a rule, they are not reduced in the cell after free radical annihilation in order to work in cycle. They only reduce the size of free radical damage.

The antioxidants which enter easily the mitochondria are the most interesting ones. Loweset al. [79] shows that the mitochondria targeted antioxidant MitoQ protects against fluoroquinolone-induced oxidative stress and mitochondrial membrane damage in human Achilles tendon cells. In cells treated with MitoQ the oxidative stress was lower and mitochondrial membrane potential was maintained.

Simonin et al. [97] report oxidative damage of collagen I to be prevented by co-administration of N-acetylcysteine (150 mg/kg) to the mice. Tsai et al. reports similar anticytotoxic effect of resveratrol [98]. Vitamins C and E belong also to this group, however, after they annihilate some radicals, they can be reduced using $NADPH_2$. Some papers point to the ability of Vitamin E to reduce the consequences of FQ-induced damage [99]. Vitamin C is presented to possess the ability to protect against lethal gamma photon irradiation in mice – strong source of OS [100]. Trace elements Zn^{2+} , $Cu^{1/2+}$, Se^{2+} , $Fe^{2/3+}$ and Mn^{2+} are co-factors of important antioxidative enzymes. Selenium supplementation is reported [101] to partially restore oxidative stress and sperm damage in FQ-treated cells.

Mn^{2+} seems to be very important, because it is a co-factor of mitochondrial SOD2 being the first barrier against O_2^- and carrying mtDNA against free radical damage. Thus, the amount of trace elements in the cell must be satisfactory. Detailed research is required for $Fe^{2/3+}$ and $Cu^{1/2+}$ in order to find if their supplementation does not increase Fenton reaction to occur. Citrate and other glycolysis inhibitors may reduce the ‘hydrogen pressure’ on ETC reducing to some degree LEC and OS.

B . Restoring the reduced mitochondrial potential may be one of important steps in restoring the proper regulatory balance in FQ-patients, however, it is not a trivial task. On the one hand, reducing OS may contribute to restoring $\Delta\Psi_m$, on the other hand, the reasons of PTP opening seem to be more composed and require advanced research. The points of interest may be re-activating HIF-1 α system, reducing intracellular and intramitochondrial Ca^{2+} concentrations, restoring membrane potential ΔV , restoring intracellular Mg^{2+} , all contributing to PTP

closing. Cyclosporine A and Metformin are postulated to be able to close PTPs [102, 103] and protect against OS [102, 104-106], thus, it is an interesting substance for possible FQAD treatment.

C . The role of uni- and bivalent cations was partially discussed in point A. Additionally, the role of Mg^{2+} and K^+ must be presented. Both ions possess high intracellular concentrations (Mg^{2+} : 20x and K^+ 40x higher than extracellular). K^+ is probably removed from the cell in the OS state by opening redox sensitive Kv1.5 channels. Mg^{2+} is strongly chelated by FQs, but, probably, it also escapes from the cells due to some unrecognized mechanisms. The supplementation must take into consideration the regulatory effect of kidney, which removes overdosed amounts of both cations to the urine. Thus, small but often doses are rather recommended in order to keep a bit higher concentrations of both ions in the blood plasma, which makes it possible to reduce the concentration gradients across the cell membranes and facilitate entering both to the cell.

D . Supporting the mitochondrial exchange (removing that destroyed ones and replication of that more healthy) is the necessary way in the case of irreversible mtDNA damage. The substance that is postulated to possess the ability to promote the mitochondrial biogenesis is pyrroloquinoline quinone (PQQ) [107, 108]. This substance is also postulated to be OS-protective [109].

E . The problem of FQ accumulation is purely present in the available research. Thus, first, it is the urgent topic for establishing if this phenomenon really, and to what extent, takes place. Removing accumulated FQs may undergo in two ways: by cytochromes P-450 in the microsomes and by different processes which can remove the molecule outside the cell. Activating the reduced cytochrome detoxification may be an important point in FQAD patients. On the other hand, ozonation has been described to be an effective method for removing first generation FQ - flumequine from the liquid water [110]. Thus, ozon therapy can be examined to be method of FQ degradation in the body.

F. Every factors presented above contribute to the disturbed gene expression which can contribute to vicious circle-like regulations causing the new regulatory balance lying far away from that optimal one. If it is this main reason of the chronic state of FQAD patients then there is a big chance to find methods for quick and effective treating of this state. However, the problem is of high complexity.

References:

1. Stephenson, A.L., et al., *Tendon Injury and Fluoroquinolone Use: A Systematic Review*. Drug Saf, 2013. **36**(9): p. 709-21.
2. Lewis, T. and J. Cook, *Fluoroquinolones and tendinopathy: a guide for athletes and sports clinicians and a systematic review of the literature*. J Athl Train, 2014. **49**(3): p. 422-7.
3. Arabyat, R.M., et al., *Fluoroquinolone-associated tendon-rupture: a summary of reports in the Food and Drug Administration's adverse event reporting system*. Expert Opin Drug Saf, 2015. **14**(11): p. 1653-60.
4. Ball, P., et al., *Comparative tolerability of the newer fluoroquinolone antibacterials*. Drug Saf, 1999. **21**(5): p. 407-21.
5. Mattappalil, A. and K.A. Mergenhagen, *Neurotoxicity with antimicrobials in the elderly: a review*. Clin Ther, 2014. **36**(11): p. 1489-1511 e4.
6. Menzies, D., N.C. Klein, and B.A. Cunha, *Trovafloxacin neurotoxicity*. Am J Med, 1999. **107**(3): p. 298-9.
7. Doussau de Bazignan, A., et al., *[Psychiatric adverse effects of fluoroquinolone: review of cases from the French pharmacologic surveillance database]*. Rev Med Interne, 2006. **27**(6): p. 448-52.
8. Kaur, K., et al., *Fluoroquinolone-related neuropsychiatric and mitochondrial toxicity: a collaborative investigation by scientists and members of a social network*. J Community Support Oncol, 2016. **14**(2): p. 54-65.
9. Moorthy, N., N. Raghavendra, and P.N. Venkatarathnamma, *Levofloxacin-induced acute psychosis*. Indian J Psychiatry, 2008. **50**(1): p. 57-8.
10. Thomas, R.J. and D.R. Reagan, *Association of a Tourette-like syndrome with ofloxacin*. Ann Pharmacother, 1996. **30**(2): p. 138-41.
11. Halkin, H., *Adverse effects of the fluoroquinolones*. Rev Infect Dis, 1988. **10 Suppl 1**: p. S258-61.
12. Francis, J.K. and E. Higgins, *Permanent Peripheral Neuropathy: A Case Report on a Rare but Serious Debilitating Side-Effect of Fluoroquinolone Administration*. J Investig Med High Impact Case Rep, 2014. **2**(3): p. 2324709614545225.
13. Hedenmalm, K. and O. Spigset, *Peripheral sensory disturbances related to treatment with fluoroquinolones*. J Antimicrob Chemother, 1996. **37**(4): p. 831-7.
14. Etminan, M., J.M. Brophy, and A. Samii, *Oral fluoroquinolone use and risk of peripheral neuropathy: a pharmacoepidemiologic study*. Neurology, 2014. **83**(14): p. 1261-3.
15. Dukewich, M., et al., *Intractable Acute Pain Related to Fluoroquinolone-Induced Peripheral Neuropathy*. J Pain Palliat Care Pharmacother, 2017: p. 1-4.
16. Tillotson, G.S., *Comment: peripheral neuropathy syndrome and fluoroquinolones*. Ann Pharmacother, 2001. **35**(12): p. 1673-4.
17. Aoun, M., et al., *Peripheral neuropathy associated with fluoroquinolones*. Lancet, 1992. **340**(8811): p. 127.
18. Ali, A.K., *Peripheral neuropathy and Guillain-Barre syndrome risks associated with exposure to systemic fluoroquinolones: a pharmacovigilance analysis*. Ann Epidemiol, 2014. **24**(4): p. 279-85.
19. Cohen, J.S., *Peripheral neuropathy associated with fluoroquinolones*. Ann Pharmacother, 2001. **35**(12): p. 1540-7.
20. Stahlmann, R. and K. Riecke, *[Well tolerated or risky? Adverse effect of quinolones]*. Pharm Unserer Zeit, 2001. **30**(5): p. 412-7.
21. Hsiao, C.J., H. Younis, and U.A. Boelsterli, *Trovafloxacin, a fluoroquinolone antibiotic with hepatotoxic potential, causes mitochondrial peroxynitrite stress in a mouse model of underlying mitochondrial dysfunction*. Chem Biol Interact, 2010. **188**(1): p. 204-13.
22. Matsubara, R., T. Kibe, and T. Nomura, *Crystalline nephropathy caused by tosufloxacin*. Pediatr Int, 2016. **58**(11): p. 1219-1221.

23. Golomb, B.A., H.J. Koslik, and A.J. Redd, *Fluoroquinolone-induced serious, persistent, multisymptom adverse effects*. BMJ Case Rep, 2015. **2015**.
24. Telfer, S.J., *Fluoroquinolone antibiotics and type 2 diabetes mellitus*. Med Hypotheses, 2014. **83**(3): p. 263-9.
25. Beckman, K.B. and B.N. Ames, *Mitochondrial aging: open questions*. Ann N Y Acad Sci, 1998. **854**: p. 118-27.
26. Chance, B., H. Sies, and A. Boveris, *Hydroperoxide metabolism in mammalian organs*. Physiol Rev, 1979. **59**(3): p. 527-605.
27. Hansford, R.G., B.A. Hogue, and V. Mildaziene, *Dependence of H₂O₂ formation by rat heart mitochondria on substrate availability and donor age*. J Bioenerg Biomembr, 1997. **29**(1): p. 89-95.
28. St-Pierre, J., et al., *Topology of superoxide production from different sites in the mitochondrial electron transport chain*. J Biol Chem, 2002. **277**(47): p. 44784-90.
29. Kudin, A.P., et al., *Characterization of superoxide-producing sites in isolated brain mitochondria*. J Biol Chem, 2004. **279**(6): p. 4127-35.
30. Brand, M.D., *Mitochondrial generation of superoxide and hydrogen peroxide as the source of mitochondrial redox signaling*. Free Radic Biol Med, 2016. **100**: p. 14-31.
31. Goncalves, R.L., et al., *Sites of superoxide and hydrogen peroxide production by muscle mitochondria assessed ex vivo under conditions mimicking rest and exercise*. J Biol Chem, 2015. **290**(1): p. 209-27.
32. Bonnet, S., et al., *A mitochondria-K⁺ channel axis is suppressed in cancer and its normalization promotes apoptosis and inhibits cancer growth*. Cancer Cell, 2007. **11**(1): p. 37-51.
33. Michelakis, E.D., et al., *Hypoxic pulmonary vasoconstriction: redox regulation of O₂-sensitive K⁺ channels by a mitochondrial O₂-sensor in resistance artery smooth muscle cells*. J Mol Cell Cardiol, 2004. **37**(6): p. 1119-36.
34. Shoshan-Barmatz, V., et al., *VDAC, a multi-functional mitochondrial protein regulating cell life and death*. Mol Aspects Med, 2010. **31**(3): p. 227-85.
35. Zorov, D.B., et al., *Regulation and pharmacology of the mitochondrial permeability transition pore*. Cardiovasc Res, 2009. **83**(2): p. 213-25.
36. Marrot, L. and J.R. Meunier, *Skin DNA photodamage and its biological consequences*. J Am Acad Dermatol, 2008. **58**(5 Suppl 2): p. S139-48.
37. Shi, Z.Y., et al., *Piperonal ciprofloxacin hydrazone induces growth arrest and apoptosis of human hepatocarcinoma SMMC-7721 cells*. Acta Pharmacol Sin, 2012. **33**(2): p. 271-8.
38. Sun, J.P., et al., *Trimethoxy-benzaldehyde levofloxacin hydrazone inducing the growth arrest and apoptosis of human hepatocarcinoma cells*. Cancer Cell Int, 2013. **13**(1): p. 67.
39. Yadav, V., et al., *Gatifloxacin induces S and G₂-phase cell cycle arrest in pancreatic cancer cells via p21/p27/p53*. PLoS One, 2012. **7**(10): p. e47796.
40. Yadav, V., et al., *Moxifloxacin and ciprofloxacin induces S-phase arrest and augments apoptotic effects of cisplatin in human pancreatic cancer cells via ERK activation*. BMC Cancer, 2015. **15**: p. 581.
41. Aranha, O., et al., *Suppression of human prostate cancer cell growth by ciprofloxacin is associated with cell cycle arrest and apoptosis*. Int J Oncol, 2003. **22**(4): p. 787-94.
42. Cencioni, C., et al., *Oxidative stress and epigenetic regulation in ageing and age-related diseases*. Int J Mol Sci, 2013. **14**(9): p. 17643-63.
43. Stahlmann, R., et al., *Ofloxacin in juvenile non-human primates and rats. Arthropathia and drug plasma concentrations*. Arch Toxicol, 1990. **64**(3): p. 193-204.
44. Maslanka, T., J.J. Jaroszewski, and M. Chrostowska, *Pathogenesis of quinolone-induced arthropathy: a review of hypotheses*. Pol J Vet Sci, 2004. **7**(4): p. 323-31.
45. Egerbacher, M., et al., *Ciprofloxacin causes cytoskeletal changes and detachment of human and rat chondrocytes in vitro*. Arch Toxicol, 2000. **73**(10-11): p. 557-63.

46. Sataric, M.V., et al., *The change of microtubule length caused by endogenous AC fields in cell.* Biosystems, 1996. **39**(2): p. 127-33.
47. Havelka, D., et al., *High-frequency electric field and radiation characteristics of cellular microtubule network.* J Theor Biol, 2011. **286**(1): p. 31-40.
48. Cifra, M., et al., *Electric field generated by axial longitudinal vibration modes of microtubule.* Biosystems, 2010. **100**(2): p. 122-31.
49. Pokorny, J., et al., *Targeting mitochondria for cancer treatment - two types of mitochondrial dysfunction.* Prague Med Rep, 2014. **115**(3-4): p. 104-19.
50. Uivarosi, V., *Metal complexes of quinolone antibiotics and their applications: an update.* Molecules, 2013. **18**(9): p. 11153-97.
51. Ma, H.H.M., F.C.K. Chiu, and R.C. Li, *Mechanistic Investigation of the Reduction in Antimicrobial Activity of Ciprofloxacin by Metal Cations.* Pharmaceutical Research, 1997. **14**(3): p. 366-370.
52. Seedher, N. and P. Agarwal, *Effect of metal ions on some pharmacologically relevant interactions involving fluoroquinolone antibiotics.* Drug Metabol Drug Interact, 2010. **25**(1-4): p. 17-24.
53. Koga, H., *High-performance liquid chromatography measurement of antimicrobial concentrations in polymorphonuclear leukocytes.* Antimicrob Agents Chemother, 1987. **31**(12): p. 1904-8.
54. Pascual, A., et al., *Uptake and intracellular activity of trovafloxacin in human phagocytes and tissue-cultured epithelial cells.* Antimicrob Agents Chemother, 1997. **41**(2): p. 274-7.
55. Andriole, V.T., *The quinolones – third edition.* 2000, San Diego California: Academic Press.
56. Badal, S., Y.F. Her, and L.J. Maher, 3rd, *Nonantibiotic Effects of Fluoroquinolones in Mammalian Cells.* J Biol Chem, 2015. **290**(36): p. 22287-97.
57. Valko, M., et al., *Redox- and non-redox-metal-induced formation of free radicals and their role in human disease.* Arch Toxicol, 2016. **90**(1): p. 1-37.
58. Shakibaei, M., et al., *Comparative evaluation of ultrastructural changes in articular cartilage of ofloxacin-treated and magnesium-deficient immature rats.* Toxicol Pathol, 1996. **24**(5): p. 580-7.
59. Egerbacher, M., et al., *Integrins mediate the effects of quinolones and magnesium deficiency on cultured rat chondrocytes.* Eur J Cell Biol, 1999. **78**(6): p. 391-7.
60. Egerbacher, M., J. Edinger, and W. Tschulenck, *Effects of enrofloxacin and ciprofloxacin hydrochloride on canine and equine chondrocytes in culture.* Am J Vet Res, 2001. **62**(5): p. 704-8.
61. Egerbacher, M., B. Wolfesberger, and C. Gabler, *In vitro evidence for effects of magnesium supplementation on quinolone-treated horse and dog chondrocytes.* Vet Pathol, 2001. **38**(2): p. 143-8.
62. Stahlmann, R., et al., *Effects of magnesium deficiency on joint cartilage in immature beagle dogs: immunohistochemistry, electron microscopy, and mineral concentrations.* Arch Toxicol, 2000. **73**(10-11): p. 573-80.
63. Stahlmann, R., et al., *Chondrotoxicity of ciprofloxacin in immature beagle dogs: immunohistochemistry, electron microscopy and drug plasma concentrations.* Arch Toxicol, 2000. **73**(10-11): p. 564-72.
64. Chui, D., L. Cheng, and A.M. Tejani, *Clinical Equivalency of Ciprofloxacin 750 mg Enterally and 400 mg Intravenously for Patients Receiving Enteral Feeding: Systematic Review.* Can J Hosp Pharm, 2009. **62**(2): p. 127-34.
65. Lomaestro, B.M. and G.R. Bailie, *Absorption interactions with fluoroquinolones. 1995 update.* Drug Saf, 1995. **12**(5): p. 314-33.
66. Marchbanks, C.R., *Drug-drug interactions with fluoroquinolones.* Pharmacotherapy, 1993. **13**(2 Pt 2): p. 235-285.
67. Muruganathan, G., et al., *Interaction study of moxifloxacin and lomefloxacin with co-administered drugs.* Pak J Pharm Sci, 2011. **24**(3): p. 339-43.

68. Polk, R.E., *Drug-drug interactions with ciprofloxacin and other fluoroquinolones*. Am J Med, 1989. **87**(5A): p. 76S-81S.
69. Radandt, J.M., C.R. Marchbanks, and M.N. Dudley, *Interactions of fluoroquinolones with other drugs: mechanisms, variability, clinical significance, and management*. Clin Infect Dis, 1992. **14**(1): p. 272-84.
70. Wright, D.H., et al., *Decreased in vitro fluoroquinolone concentrations after admixture with an enteral feeding formulation*. JPEN J Parenter Enteral Nutr, 2000. **24**(1): p. 42-8.
71. Qin, P. and R. Liu, *Oxidative stress response of two fluoroquinolones with catalase and erythrocytes: a combined molecular and cellular study*. J Hazard Mater, 2013. **252-253**: p. 321-9.
72. Yu, C.H., et al., *Effect of Danofloxacin on Reactive Oxygen Species Production, Lipid Peroxidation and Antioxidant Enzyme Activities in Kidney Tubular Epithelial Cell Line, LLC-PK1*. Basic Clin Pharmacol Toxicol, 2013. **113**(6): p. 377-84.
73. Pouzaud, F., et al., *In vitro discrimination of fluoroquinolones toxicity on tendon cells: involvement of oxidative stress*. J Pharmacol Exp Ther, 2004. **308**(1): p. 394-402.
74. Pouzaud, F., et al., *Age-dependent effects on redox status, oxidative stress, mitochondrial activity and toxicity induced by fluoroquinolones on primary cultures of rabbit tendon cells*. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology, 2006. **143**(2): p. 232-241.
75. Kumbhar, G.B., A.M. Khan, and S. Rampal, *Evaluation of gatifloxacin for its potential to induce antioxidant imbalance and retinopathy in rabbits*. Hum Exp Toxicol, 2015. **34**(4): p. 372-9.
76. Talla, V. and P. Veerareddy, *Oxidative stress induced by fluoroquinolones on treatment for complicated urinary tract infections in Indian patients*. J Young Pharm, 2011. **3**(4): p. 304-9.
77. Liu, B., et al., *Cytotoxic effects and apoptosis induction of enrofloxacin in hepatic cell line of grass carp (Ctenopharyngodon idellus)*. Fish Shellfish Immunol, 2015. **47**(2): p. 639-44.
78. Li, H.T., S.Y. Zhu, and H.S. Pei, *[The effect of moxifloxacin on apoptosis of airway smooth muscle cells and mitochondria membrane potential]*. Zhonghua Jie He He Hu Xi Za Zhi, 2011. **34**(9): p. 684-7.
79. Lowes, D.A., et al., *The mitochondria targeted antioxidant MitoQ protects against fluoroquinolone-induced oxidative stress and mitochondrial membrane damage in human Achilles tendon cells*. Free Radic Res, 2009. **43**(4): p. 323-8.
80. Madesh, M. and G. Hajnoczky, *VDAC-dependent permeabilization of the outer mitochondrial membrane by superoxide induces rapid and massive cytochrome c release*. J Cell Biol, 2001. **155**(6): p. 1003-15.
81. Simamura, E., et al., *Furanonaphthoquinones cause apoptosis of cancer cells by inducing the production of reactive oxygen species by the mitochondrial voltage-dependent anion channel*. Cancer Biol Ther, 2006. **5**(11): p. 1523-9.
82. Akahane, K., et al., *Structure-epileptogenicity relationship of quinolones with special reference to their interaction with gamma-aminobutyric acid receptor sites*. Antimicrob Agents Chemother, 1989. **33**(10): p. 1704-8.
83. Akahane, K., et al., *Possible intermolecular interaction between quinolones and biphenylacetic acid inhibits gamma-aminobutyric acid receptor sites*. Antimicrob Agents Chemother, 1994. **38**(10): p. 2323-9.
84. Divakaruni, A.S. and M.D. Brand, *The regulation and physiology of mitochondrial proton leak*. Physiology (Bethesda), 2011. **26**(3): p. 192-205.
85. Ferrandiz, M.J. and A.G. de la Campa, *The fluoroquinolone levofloxacin triggers the transcriptional activation of iron transport genes that contribute to cell death in Streptococcus pneumoniae*. Antimicrob Agents Chemother, 2014. **58**(1): p. 247-57.
86. Ferrandiz, M.J., et al., *Reactive Oxygen Species Contribute to the Bactericidal Effects of the Fluoroquinolone Moxifloxacin in Streptococcus pneumoniae*. Antimicrob Agents Chemother, 2015. **60**(1): p. 409-17.

87. Gong, Y., et al., *Partial degradation of levofloxacin for biodegradability improvement by electro-Fenton process using an activated carbon fiber felt cathode*. J Hazard Mater, 2016. **304**: p. 320-8.
88. Michael, I., et al., *Solar Fenton and solar TiO₂ catalytic treatment of ofloxacin in secondary treated effluents: evaluation of operational and kinetic parameters*. Water Res, 2010. **44**(18): p. 5450-62.
89. Fox, A.J., et al., *Fluoroquinolones impair tendon healing in a rat rotator cuff repair model: a preliminary study*. Am J Sports Med, 2014. **42**(12): p. 2851-9.
90. Liang, X., et al., *Effects of norfloxacin on hepatic genes expression of P450 isoforms (CYP1A and CYP3A), GST and P-glycoprotein (P-gp) in Swordtail fish (Xiphophorus Helleri)*. Ecotoxicology, 2015. **24**(7-8): p. 1566-73.
91. Outman, W.R. and C.H. Nightingale, *Metabolism and the fluoroquinolones*. Am J Med, 1989. **87**(6C): p. 37S-42S.
92. Shlosberg, A., et al., *The inhibitory effects of the fluoroquinolone antimicrobials norfloxacin and enrofloxacin on hepatic microsomal cytochrome P-450 monooxygenases in broiler chickens*. Drug Metabol Drug Interact, 1997. **14**(2): p. 109-22.
93. Granfors, M.T., et al., *Ciprofloxacin greatly increases concentrations and hypotensive effect of tizanidine by inhibiting its cytochrome P450 1A2-mediated presystemic metabolism*. Clin Pharmacol Ther, 2004. **76**(6): p. 598-606.
94. Regmi, N.L., et al., *Lack of inhibitory effects of several fluoroquinolones on cytochrome P-450 3A activities at clinical dosage in dogs*. J Vet Pharmacol Ther, 2007. **30**(1): p. 37-42.
95. Regmi, N.L., et al., *Inhibitory effect of several fluoroquinolones on hepatic microsomal cytochrome P-450 1A activities in dogs*. J Vet Pharmacol Ther, 2005. **28**(6): p. 553-7.
96. Brand, M.D., et al., *Suppressors of Superoxide-H₂O₂ Production at Site IQ of Mitochondrial Complex I Protect against Stem Cell Hyperplasia and Ischemia-Reperfusion Injury*. Cell Metab, 2016. **24**(4): p. 582-592.
97. Simonin, M.A., et al., *Pefloxacin-induced achilles tendon toxicity in rodents: biochemical changes in proteoglycan synthesis and oxidative damage to collagen*. Antimicrob Agents Chemother, 2000. **44**(4): p. 867-72.
98. Tsai, T.Y., et al., *The effect of resveratrol on protecting corneal epithelial cells from cytotoxicity caused by moxifloxacin and benzalkonium chloride*. Invest Ophthalmol Vis Sci, 2015. **56**(3): p. 1575-84.
99. Gurbay, A., et al., *Ciprofloxacin-induced DNA damage in primary culture of rat astrocytes and protection by Vitamin E*. Neurotoxicology, 2006. **27**(1): p. 6-10.
100. Mortazavi, S.M., et al., *A Comparative Study on the Life-Saving Radioprotective Effects of Vitamins A, E, C and Over-the-Counter Multivitamins*. J Biomed Phys Eng, 2015. **5**(2): p. 59-66.
101. Rungsung, S., et al., *Evaluation of ameliorative potential of supranutritional selenium on enrofloxacin-induced testicular toxicity*. Chem Biol Interact, 2016. **252**: p. 87-92.
102. Detaille, D., et al., *Metformin prevents high-glucose-induced endothelial cell death through a mitochondrial permeability transition-dependent process*. Diabetes, 2005. **54**(7): p. 2179-87.
103. El-Mir, M.Y., et al., *Neuroprotective role of antidiabetic drug metformin against apoptotic cell death in primary cortical neurons*. J Mol Neurosci, 2008. **34**(1): p. 77-87.
104. Lee, J.Y., et al., *Protective effect of metformin on gentamicin-induced vestibulotoxicity in rat primary cell culture*. Clin Exp Otorhinolaryngol, 2014. **7**(4): p. 286-94.
105. Salman, Z.K., et al., *The combined effect of metformin and L-cysteine on inflammation, oxidative stress and insulin resistance in streptozotocin-induced type 2 diabetes in rats*. Eur J Pharmacol, 2013. **714**(1-3): p. 448-55.
106. Morales, A.I., et al., *Metformin prevents experimental gentamicin-induced nephropathy by a mitochondria-dependent pathway*. Kidney Int, 2010. **77**(10): p. 861-9.
107. Chowanadisai, W., et al., *Pyrroloquinoline quinone stimulates mitochondrial biogenesis through cAMP response element-binding protein phosphorylation and increased PGC-1alpha expression*. J Biol Chem, 2010. **285**(1): p. 142-52.

108. Stites, T., et al., *Pyroloquinoline quinone modulates mitochondrial quantity and function in mice*. J Nutr, 2006. **136**(2): p. 390-6.
109. Huang, Y., N. Chen, and D. Miao, *Biological effects of pyroloquinoline quinone on liver damage in Bmi-1 knockout mice*. Exp Ther Med, 2015. **10**(2): p. 451-458.
110. Feng, M., et al., *Fast removal of the antibiotic flumequine from aqueous solution by ozonation: Influencing factors, reaction pathways, and toxicity evaluation*. Sci Total Environ, 2016. **541**: p. 167-75.